

Reply to Fedson

TO THE EDITOR—We appreciate the interest in our recent article [1] on biomarkers in Ebola virus disease (EVD) and the editorial comments provided by Fedson. In his editorial, Fedson points out that the evidence of endothelial dysfunction that we reported in patients affected by EVD is also commonly observed in other forms of sepsis. He brings attention to a mouse study that indicated that selective inhibition of nuclear factor κ B signaling in endothelial cells in the context of bacterial sepsis provided protection from disease [2], suggesting that downregulation of the local inflammatory response at the endothelial cell level could be protective to patients. Additionally, he comments on a small clinical trial of statins in patients with sepsis; this study showed decreased progression to severe sepsis in statin-treated patients [3]. This result is attributed to the role of statins in stabilizing the endothelium and in its local antiinflammatory effects on endothelial cells. Finally, Fedson suggests the use of statins as adjunctive therapy in managing patients who have EVD.

We agree that statins should be considered as adjunctive therapy for EVD. In fact, we suggested the use of statins in patients with EVD in a subsequent study, in which we examined the differences in biomarkers and clinical outcomes between adult and pediatric patients with EVD [4]. In that study, we observed direct evidence of endothelial dysfunction, demonstrated by elevated levels of soluble intercellular adhesion molecule and soluble vascular cell adhesion molecule

in pediatric patients who had fatal outcomes, and normal levels of these factors in pediatric patients who survived infection.

No treatment is without risk, however, and the benefits and risks of a given intervention or medication must always be considered. The risks of using statins include rhabdomyolysis, leading to acute renal failure, and many potential drug interactions, since statins are metabolized by the cytochrome P450 system. Also, patients with known elevation of hepatic transaminase levels are at increased risk of side effects. Since Ebola virus is known to be hepatotropic and severe disease is associated with elevated hepatic transaminase levels, patients would need to be carefully monitored if statins were initiated. This type of intensive monitoring could be very difficult to achieve in the areas of the world affected by EVD. On the other hand, use of statins in patients with EVD could stabilize the endothelium by decreasing local inflammation and, thus, lessen the risk of the patient developing shock. Additionally, although only 30%–40% of patients develop hemorrhagic manifestations of EVD and because the presence of hemorrhage does not correlate with death, statin use might also decrease the incidence of hemorrhage by affecting components of the coagulation pathway.

Unanswered questions include the dose, frequency, duration, and choice of statin to use in patients with EVD. Presumably, initiating statin treatment within 24 hours of the onset of symptoms would be best, although given the severity of untreated EVD, initiation of treatment at any point in the disease course would be reasonable. In the ASEPSIS trial, 40 mg of atorvastatin was administered daily for the duration of hospitalization, up to 28 days. Patients who survive EVD tend to show recovery within 2 weeks of symptom onset, so this duration of therapy would theoretically be sufficient. Typically, a potentially useful drug would be evaluated in several animal models of disease long before its use in humans is even considered. Then, a rigorous, double-

blinded, placebo-controlled trial would be conducted to evaluate that drug in humans. However, we are in the midst of the largest EVD outbreak ever recorded, and it shows no signs of abating. Therefore, the individual clinician is left with the decision to use a drug off-label that might be beneficial, while weighing the possible side effects of that drug. These decisions should be made very carefully, in consultation with the patient, and with full disclosure of the possible risks and benefits.

Note

Potential conflicts of interest. Both authors: No reported conflicts.

All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

Anita K. McElroy^{1,2} and Christina F. Spiropoulou²

¹Emory Pediatric Infectious Disease, and ²US Centers for Disease Control and Prevention, Viral Special Pathogens Branch, Atlanta, Georgia

References

1. McElroy AK, Erickson BR, Flietstra TD, et al. Ebola hemorrhagic Fever: novel biomarker correlates of clinical outcome. *J Infect Dis* **2014**; 210:558–66.
2. Xu H, Ye X, Steinberg H, Liu SF. Selective blockade of endothelial NF-kappaB pathway differentially affects systemic inflammation and multiple organ dysfunction and injury in septic mice. *J Pathol* **2010**; 220: 490–8.
3. Patel JM, Snaith C, Thickett DR, et al. Randomized double-blind placebo-controlled trial of 40 mg/day of atorvastatin in reducing the severity of sepsis in ward patients (ASEP-SIS Trial). *Crit Care* **2012**; 16:R231.
4. McElroy AK, Erickson BR, Flietstra TD, et al. Biomarker correlates of survival in pediatric patients with Ebola virus disease. *Emerg Infect Dis* **2014**. In press. http://wwwnc.cdc.gov/eid/article/20/10/14-0430_article. Accessed 13 August 2014.

Received 14 August 2014; accepted 18 August 2014; electronically published 25 August 2014.

Correspondence: Anita K. McElroy, MD, PhD, Division of Infectious Disease, Department of Pediatrics, Emory University, 2015 Uppergate Drive, Suite 500, Atlanta, GA (akmcelr@emory.edu).

The Journal of Infectious Diseases® 2015;211:662–3

Published by Oxford University Press on behalf of the Infectious Diseases Society of America 2014. This work is written by (a) US Government employee(s) and is in the public domain in the US. DOI: 10.1093/infdis/jiu475